

Complications of Diagnostic Intra-arterial Cerebral Angiography in Ischemic Cerebrovascular Diseases

Kyung Yul Lee, M.D., Seo Hyun Kim, M.D., Pyeong Ho Yoon, M.D.*, Dong Ik Kim, M.D.*, Byung In Lee, M.D., Seung Min Kim, M.D., Soo Chul Park, M.D., Young Ho Sohn, M.D., Ji Hoe Heo, M.D.

Department of Neurology and Diagnostic Radiology, Yonsei University College of Medicine*

Background : Intra-arterial cerebral angiography, which is a prerequisite for carotid endarterectomy and angioplasty, carries some risks but provides the best visualization of the cerebral vasculatures. We attempted to examine the incidence of complications associated with cerebral angiography in patients with ischemic stroke. **Methods** : We retrospectively reviewed the medical records of patients with ischemic stroke or transient ischemic attack (TIA) who underwent the digital subtraction cerebral angiography. Four hundred nineteen procedures were performed between October 1994 and August 1999. The systemic, local, and neurologic complications were evaluated. The neurologic complications were defined as occurrences of any new focal neurologic deficits or progressions of the preexisting neurologic deficits during or within 24 hours after the procedure. **Results** : There were 5 systemic (1.2%), 17 local (4.1%), and 10 neurologic (2.4%) complications. The neurologic complications were reversible within 7 days in 6 (1.4%) and were persistent after 7 days in 4 (1.0%). Six out of 10 patients with neurologic complications had previous stroke or TIA. The angiographic studies revealed the stenosis or obstruction of the relevant arteries in 7 patients. **Conclusions** : Cerebral angiography in patients with ischemic stroke was associated with 1.4 % reversible and 1.0% persistent neurologic complications, all of which developed after the angiographic procedure. The history of previous stroke or TIA and the presence of severe stenosis or occlusion of the symptomatic arteries may carry a high risk of neurologic complications.

J Korean Neurol Assoc 19(4):354~358, 2001

Key Words : Cerebral ischemia, Cerebral angiography, Complication

가

가 가

NASCET¹

가

ECST²

가

가

가

Manuscript received March 29, 2001

Accepted in final form May 17, 2001

* Address for correspondence

Ji Hoe Heo, M.D.

Department of Neurology, Yonsei University College of Medicine, Shinchon-dong 134, Seodaemun-gu, Seoul, 120-752, Korea

Tel : +82-2-361-5467 Fax : +82-2-393-0705

E-mail : jhheo@yumc.yonsei.ac.kr

* 1995 1996 (HMP-96-M-2-0015)

24

1994 10 1999 8

24

(transient), 1
(reversible), 1
(persistent)가
(persis-
tent)

Yonsei Stroke Registry,

419 가 289 , 가 130

57.2 ± 11.1 (20 ~ 82)

(TIA) 가 42 , 가 377

SPSS for Windows, ver 9.0

t-test

105 ,

Chi-

square test

57 , 132 ,

가 75 , 8

digital subtraction angiography

Seldinger

5 (1.2%)

419

. 4

1

4 vessel-study

3

24

. 4

1

24

(cardiac arrhythmia)

Table 1. Summary of neurologic complications after digital subtraction cerebral angiography

Patient	Age/Sex	Complications	Interval (hr)	History of previous stroke or TIA	Angiographic findings
With persistent complications					
1	44/M	Left sided weakness developed progressive	22		Right cervical ICA occlusion left cavernous ICA, VA severe stenosis
2	78/M	Left sided weakness aggravated	2	Yes	Right cervical ICA occlusion
3	63/M	Right sided weakness developed progressive	24	Yes	Right proximal MCA moderate stenosis left cervical ICA severe stenosis
4	67/M	Right sided weakness developed	12		Bilateral cavernous ICA mild stenosis
With reversible complications					
5	55/F	Aphasia aggravated	4		Left cavernous ICA severe stenosis
6	64/F	Right sided weakness aggravated	1	Yes	Normal
7	54/M	Diplopia and left sided clumsiness developed	17	Yes	Left ICA total occlusion
8	65/M	Dysarthria and right sided weakness developed	14		Right cavernous ICA severe stenosis left MCA moderate stenosis
9	61/F	Sensory aphasia developed	3	Yes	Left MCA moderate stenosis bilateral PCA moderate stenosis
10	56/F	Right sided weakness developed	8	Yes	Left MCA moderate stenosis

TIA: transient ischemic attack, ICA: internal carotid artery, MCA: middle cerebral artery, VA: vertebral artery

PCA: posterior cerebral artery, mild: 30-50 % stenosis, moderate: 51-70% stenosis, severe: 71-99 % stenosis

Interval means the time lapse to the onset of neurologic signs after the end of the angiographic procedure

Table 2. Neurologic complications associated with cerebral angiography in the literatures

Studies	Number of patients	Neurologic complications (%)		Note
		Total	Permanent	
Dion et al ³	724	1.3	0.1	
Hankey et al ⁴	382	2.6	1.3	S
Grzyska et al ⁵	1095	0.54	0.09	D
Waugh et al ⁶	939	0.9	0.3	D
Warnock et al ⁷	395	3.89	0.52	D,S
Heiserman et al ⁸	688	1.0	0.5	
Leffers et al ⁹	454	2.3	0.4	D
Choi et al ¹⁰	171	4.6	1.8	D
Present study	419	2.4	1.0	D,S

D: Digital subtraction cerebral angiographic study, S: only patients with ischemic cerebrovascular diseases

17 (4.1%) (p<0.01).
2 7 (1-3, 5, 8-10)
가 (pseudoaneurysm)가
15 , 1 (4)
17 2
가 가 2 24
가 (arterial 1 (1)
fistula) , 1 (2)
10 (2.4%)
60.7 419
(44~78)
(57.1)
가
3 7 0.5~4.7%
(Table 2).³⁻¹⁰
24 (1~24 , 10.7
) 3
4 가
7 5 12
(Table 1). 10 6 (1.4
%) 7 가
4 (1.0%) 7 가
2 가 3,5,8,12
(progressive stroke) 2.4%
24 1.0%
5
1 1.0% 가
2.5% .¹¹
409
47 Warnock

- bral angiography: A retrospective study of complication rate and patient risk factors. *Acta Radiol* 2000;41:204-210.
10. Choi JY, Han SR, Kim YI. Complications of cerebral angiography in the patients with cerebrovascular diseases. *Korean J Stroke* 1999;1:197-200.
 11. Cooperative Study between the ASNR, ASITN, and the SCVIR. Quality improvement guidelines for adult diagnostic neuroangiography. *AJNR* 2000;21:146-150.
 12. Hankey GJ, Warlow CP, Sellar RJ. Cerebral angiographic risk in mild cerebrovascular disease. *Stroke* 1990;21:209-222.
 13. Britt PM, Heiserman JE, Snider RM, Shill HA, Bird CR, Wallace RC. Incidence of postangiographic abnormalities revealed by diffusion-weighted MR imaging. *AJNR* 2000;21:55-59.
 14. Bendszus M, Koltzenburg M, Burger R, Warmuth-Metz M, Hofmann E, Solymosi L. Silent embolism in diagnostic cerebral angiography and neurointerventional procedures: A prospective study. *Lancet* 1999;354:1594-1597.
 15. Dagirmanjian A, Davis DA, Rothfus WE, Deeb ZL, Goldberg AL. Silent cerebral microemboli occurring during carotid angiography: Frequency as determined with Doppler sonography. *AJR* 1993;161:1037-1040.
 16. Gerraty RP, Bowser DN, Infeld B, Mitchell PJ, Davis SM. Microemboli during carotid angiography: Association with stroke risk factors or subsequent magnetic resonance imaging changes? *Stroke* 1996;27:1543-1547.
 17. Abeyama K, Oh S, Kawano K, Nakajima T, Soejima Y, Nakano K, et al. Nonionic contrast agents produce thrombotic effect by inducing adhesion of leukocytes on human endothelium. *Biochem Biophys Res Commun* 1995;212:776-783.
 18. Barstad RM, Buchmann MS, Hamers MJ, Orning L, Orvim U, Stormorken H, et al. Effects of ionic and non-ionic contrast media on endothelium and on arterial thrombus formation. *Acta Radiol* 1996;37:954-961.
 19. Chronos NA, Goodall AH, Wilson DJ, Sigwart U, Buller NP. Profound platelet degranulation is an important side effect of some types of contrast media used in interventional cardiology. *Circulation* 1993;88:2035-2044.
 20. Gabrielsen TO. Neurologic complications of cerebral angiography. *AJNR* 1994;15:1408-1411.